


Problem set 2: Metropolis Criterion, Ising Model

1. a. You have generated 100000 configurations of a Lennard-Jones system using (1) simple sampling, (2) importance sampling. How do you calculate the average energy?
 - b. Verify that the Metropolis Criterion enforces detailed-balance. (Distinguish between $\Delta E < 0$ und $\Delta E > 0$.)
 - c. The Glauber algorithm has the following acceptance rule: $W_{ij} = (1 - \tanh(\beta(E_j - E_i)/2))$. Show that this algorithm enforces detailed balance, too. (Hint: $\tanh(x) = (e^x - e^{-x}) / (e^x + e^{-x})$.)
 - d. Formulate the Metropolis Criterion for a system of hard discs in the canonical ensemble.

2. Consider a d-dimensional system of spins, which can point up ($s_i = +1$) or down ($s_i = -1$). Only nearest neighbors interact („Ising-Model“). We would like to write a Monte Carlo program with which we can determine the statistical properties of such a system.



$$E = -\frac{1}{2} \sum_{i,j=1}^n s_i s_j$$



- a. How many interactions need to be calculated after a single spin flip in $d=1,2,3$ dimensions?

b. Write down which steps need to be implemented to simulate the Ising model.

c. Which configurations would you expect at high and at low temperatures. Distinguish between $d=1$ and $d=2,3$.

d. If you have access to a computer: Verify part c for $d=2$:

(<http://bartok.ucsc.edu/peter/java/ising/keep>

[/ising.html](#)) What happens at $T=2.629$?

Homework: Read the (German) handout on Monte Carlo simulations.

Picture credits: www.moviewallpapers.net, oscar.cacr.caltech.edu/Hrothgar/Ising/Ising1.JPG